

## PUBLIKATIONEN

- Einfluss eines kolonie-stimulierenden Faktors der Hämatopoese auf die antikörperabhängige zelluläre Zytotoxizität von menschlichen neutrophilen Granulozyten. Dissertationsarbeit. **S. Oez**, 1985.
- Human pluripotent hematopoietic colony stimulating factor: Activities on human and murine cells. E. Platzer, **S. Oez**, K. Welte, A. Sendler, J. L. Gabilove, R. Mertelsmann, M. A. S. Moore, and J. R. Kalden. Immunobiol. 172:185-193, 1986.
- Anti-tumor efficacy of interleukin-2-activated killer cells in human neuroblastoma ex-vivo. J. Atzpodien, S. C. Gulati, J. H. Kwon, B. H. Kushner, C. Shimazaki, C. Bühner, **S. Oez**, J. E. Kolitz, K. Welte, and B. D. Clarkson. Exp. Cell Biol. 56:236-244, 1988.
- Ewing's sarcoma: Ex-vivo sensitivity towards natural (NK) and lymphokine-activated (LAK) killing. J. Atzpodien, S. C. Gulati, C. Shimazaki, C. Bühner, **S. Oez**, J. H. Kwon, J. E. Kolitz, and B. D. Clarkson. Oncology 45:437-443, 1988.
- Tumor necrosis factor regulates antibody dependent cellular cytotoxicity (ADCC) of human polymorph nuclear cells. **S. Oez**, M. Hadam, C. Bühner, E. Platzer, J. Atzpodien, and K. Welte. Cancer Res. Clin. Oncol. 114:146-146, 1988.
- A Quantitative Colorimetric Method to Assess the Functional State of Human Polymorphnuclear Leukocytes. **S. Oez**, E. Platzer, and K. Welte. Blut 60:97-102, 1990.
- Establishment and characterization of a granulocyte-macrophage colony-stimulating factor-dependent human myeloid cell line. **S. Oez**, H. Tittelbach, R. Fahsold, R. Schaeztl, and J. R. Kalden. Blood 76:578-582, 1990.
- A highly sensitive quantitative bioassay for human granulocyte- macrophage colony-stimulating factor. **S. Oez**, J. Atzpodien, C. Bühner, and J. R. Kalden. Exp. Hematol. 18:1108-1111, 1990.
- A simple assay for quantifying the inducible adherence of neutrophils. **S. Oez**, K. Welte, E. Platzer, and J. R. Kalden. Immunobiol. 180:308-315, 1990.
- Clonal growth of functionally normal and deficient neutrophils from the bone marrow of a patient with variant chronic granulomatous disease. **S. Oez**, J. Birkmann, and J. R. Kalden. Ann. Hematol. 66:21-25, 1993.
- Granulocyte-macrophage colony stimulating factor (GM-CSF) reduces the density of stem cell factor receptors (c-kit oncogene product) on a GM-CSF dependent human myeloid cell line. **S. Oez**, G. Hofmann-Wackersreuther, J. Birkmann, M. Smetak, K. Welte, and W. M. Gallmeier. Eur. Cytokine Netw. 4:293-297, 1993.
- Regulation of the density of the stem cell factor receptor (c-kit) by tumor necrosis factor on a human myeloid cell line. **S. Oez**, J. Birkmann, M. Smetak, S. Corbacioglu, G. Hofmann-Wackersreuther, K. Welte, and W. M. Gallmeier. Eur. Cytokine Netw. 4:439-445, 1993.

- LW/SO cell line: a tool for studying the phenotypical characterization and commitment of hematopoietic stem cells. **S. Oez**, U. Trautmann, M. Smetak, J. Birkmann, S. Al Salameh, E. Gebhart, and W. M. Gallmeier. *Ann. Hematol* 72:307-316, 1996.
- Effects of recombinant human thrombopoietin alone and in combination with erythropoietin and early-acting cytokines on human mobilized purified CD34+ progenitor cells cultured in serum-depleted medium. J. Birkmann, **S. Oez**, M. Smetak, G. Kaiser, H. Kappauf, and W. M. Gallmeier. *STEM Cells* 1997;15:18-32.